

## **ANALYSIS OF SWINE MOVEMENTS IN A PROVINCE IN NORTHERN VIETNAM AND APPLICATION IN THE DESIGN OF SURVEILLANCE STRATEGIES FOR INFECTIOUS DISEASES<sup>2</sup>**

**E. Baudon<sup>1,2</sup>, G. Fournie<sup>3</sup>, D. T. Hiep<sup>4</sup>, T. T. H. Pham<sup>2</sup>, R. Duboz<sup>2</sup>, M. Ge ly<sup>2</sup>,  
M. Peiris<sup>1</sup>, B. J. Cowling<sup>1</sup>, V. D. Ton<sup>4\*</sup> and M. Peyre<sup>2</sup>**

<sup>1</sup>*School of Public Health, The University of Hong Kong, Pokfulam, Hong Kong Special Administrative Region, China;* <sup>2</sup>*Animal and Integrated Risk Management Research Unit (AGIRs), French Agricultural Research Center for International Development (CIRAD), Montpellier, France;*

<sup>3</sup>*Veterinary Epidemiology and Public Health Group, Production and Population Health Department, Royal Veterinary College, North Mymms Hatfield, Hertfordshire, UK;*

<sup>4</sup>*Vietnam National University of Agriculture*

Email\*: [vdton@vnua.edu.vn](mailto:vdton@vnua.edu.vn)

### **ABSTRACT**

While swine production is rapidly growing in South-East Asia, the structure of the swine industry and the dynamic of pig movements have not been well-studied. However, this knowledge is a prerequisite for understanding the dynamic of disease transmission in swine populations and designing cost-effective surveillance strategies for infectious diseases. In this study, we assessed the farming and trading practices in the Vietnamese swine familial farming sector, which accounts for most pigs in Vietnam, and for which disease surveillance is a major challenge. Farmers from two communes of a Red River Delta Province (northern Vietnam) were interviewed, along with traders involved in pig transactions. Major differences in the trade structure were observed between the two communes. One commune had mainly transversal trades that is between farms of equivalent sizes, whereas the other had pyramidal trades that is from larger to smaller farms. Companies and large familial farrow-to-finish farms were likely to act as major sources of disease spread through pig sales, demonstrating their importance for disease control. Familial fattening farms with high pig purchases were at greater risk of disease introduction and should be targeted for disease detection as part of a risk-based surveillance. In contrast, many other familial farms were isolated or weakly connected to the swine trade network limiting their relevance for surveillance activities. However, some of these farms used boar hiring for breeding, increasing the risk of disease spread. Most familial farms were slaughtering pigs at the farm or in small local slaughterhouses, making the surveillance at the slaughterhouse inefficient. In terms of spatial distribution of the trades, the results suggested that northern provinces were highly connected and showed some connection with central and southern provinces. These results are useful to develop risk-based surveillance protocols for disease detection in the swine familial sector and to make recommendations for disease control.

**Keywords:** Network analysis, swine movements, Infectious disease, disease surveillance, South-East Asia, Vietnam.

### **Phân tích các dòng chu chuyển của lợn tại một tỉnh ở miền Bắc Việt Nam và áp dụng trong xây dựng các chiến lược giám sát bệnh truyền nhiễm**

### **TÓM TẮT**

Trong khi sản lượng lợn đang tăng trưởng nhanh chóng ở khu vực Đông Nam Á, thì cơ cấu ngành chăn nuôi lợn và động thái của các dòng chu chuyển lợn chưa được nghiên cứu nhiều. Song, sự hiểu biết về vấn đề này là một điều kiện tiên quyết cho việc lý giải động thái lây truyền bệnh trong các quần thể lợn và thiết kế các chiến lược giám sát chi phí-hiệu quả đối với các bệnh truyền nhiễm. Trong nghiên cứu này, chúng tôi đánh giá việc chăn nuôi và

<sup>2</sup> The article was published in Transboundary and Emerging Diseases. Doi: 10.1111/tbed.12380.

buôn bán lợn trong khu vực nông hộ ở Việt nam, nơi đóng vai trò quan trọng trong sản xuất và thương mại lợn ở Việt Nam, và việc giám sát dịch bệnh là một thách thức lớn. Các hộ gia đình được lựa chọn phỏng vấn từ hai xã của một tỉnh đồng bằng sông Hồng (phía Bắc Việt Nam), cùng với các thương lái tham gia vào quá trình mua bán lợn. Có sự khác biệt lớn trong cơ cấu dao dịch đã quan sát được ở hai xã. Một xã thì việc buôn bán lợn chủ yếu diễn ra theo chiều ngang đó là sự mua bán giữa các nông hộ có quy mô tương đương nhau, trong khi đó ở xã kia thì giao dịch buôn bán lại theo hình tháp đó là từ trang trại lớn tới các trang trại nhỏ. Các công ty và trang trại chăn nuôi lợn sản xuất lợn con để bán có thể là nguồn lây nhiễm bệnh chính và cần phải được kiểm soát dịch bệnh. Các hộ nuôi lợn thịt có mức mua bán lợn cao cũng có mức nguy cơ lây nhiễm cao và cần được kiểm soát. Trái lại, nhiều hộ chăn nuôi khác khá độc lập hoặc ít mua bán trao đổi có thể không đòi hỏi sự giám sát cao. Tuy nhiên, một số các hộ này nuôi lợn đực giống để kinh doanh sẽ có nguy cơ lây lan bệnh cao hơn. Hầu hết các nông hộ đã và đang giết mổ lợn tại gia đình hoặc trong các lò giết mổ nhỏ tại địa phương dẫn đến việc chỉ giám sát tại các lò mổ không hiệu quả. Liên quan đến phân bố không gian thương mại, các kết quả cho thấy rằng các tỉnh phía Bắc có giao dịch nhiều với một số tỉnh miền Trung và miền Nam. Những kết quả này là rất hữu ích cho việc xây dựng các chương trình giám sát phát hiện bệnh dịch trong khu vực chăn nuôi lợn nông hộ gia và đưa ra các khuyến nghị để kiểm soát dịch bệnh.

Từ khóa: Phân tích mạng lưới, chu chuyển lợn, bệnh truyền nhiễm, giám sát dịch bệnh, Đông Nam Á, Việt Nam.

## 1. INTRODUCTION

South-East Asia is considered as a hotspot for the emergence of zoonotic infectious diseases mostly due to anthro- pogenic factors (Jones et al., 2008; Coker et al., 2011). Swine is a host, mixing vessel or reservoir for many viral (e.g. influenza, Japanese encephalitis, Nipah virus, hepatitis E) and bacterial (e.g. *Streptococcus suis*) zoonoses which are major public health concerns in South-East Asia (Smith et al., 2011; Jones et al., 2013). The 2009 H1N1 influenza pandemic confirmed the importance of swine in the generation process of new influenza reassortants from avian, human and swine influenza viruses (SIVs). It also stressed the need for having an effective systematic surveillance of influenza viruses circulating in pigs at the global level (Vijaykrishna et al., 2011). The pig and poultry populations have at least doubled since 1990 in many South-East Asian countries, including Vietnam (Coker et al., 2011). Vietnam is the world's 13th most populous country with 88.8 million inhabitants (GSO, 2012), 4th largest pig producer with 26.2 million heads and 10th largest poultry producer with 315.0 million heads (FAO, 2014). However, no sustainable surveillance programme focusing on zoonotic diseases in swine such as SIVs has been implemented so far; only shortterm research projects were carried out. Also, only a limited passive surveillance system is established for very contagious swine diseases such as porcine reproductive and respiratory syndrome (PRRS).

The knowledge of the structure and the dynamic of the pig value chain is a prerequisite for understanding infectious disease transmission dynamics and for the development of cost-effective surveillance systems. In Vietnam, the structure of the swine industry and its implication for disease surveillance have not been extensively studied. There are three main different types of farms: familial farms (Hộ chăn nuôi), companies (Doanh nghiệp) and state-owned farms (Trại nhà nước). Familial farms are held by households and often characterized as small (Nông hộ), medium (Gia trại) or large farms (Trang trại); the actual size of each farm category may differ across the locations. Company farms are very large farms with a different legal status and ownership. They are owned by foreign or Vietnamese companies and usually under the management of several people; some companies own many farms. They also practice contract farming (Gia công) by providing financial and technical support to farmers. These farms are usually larger than the average familial farms. The company farms are mainly located around Hanoi and Ho Chi Minh City. Finally, state-owned farms are large farms which often keep exotic breed great grandparents and grandparents. The familial small and medium holders (described as 1-100 fattening pigs or 5-20 sows) consisted of 84% of national pig herds in 2006, with a decline in the number of smallholdings and an increase in the number of medium sized holdings since 1999 (Fisher and

Gordon, 2008). Indeed, the government promotes intensification of pig production, and as the smallholdings are often less robust to disease outbreaks, many small farmers stopped raising pigs, while other familial holdings converted to large-scale pig production. In most familial farms and in some companies, the biosecurity level is low, promoting inter-species transmission of diseases, such as influenza, and within and between-herd transmission of swine diseases of economic importance, such as PRRS, classical swine fever and foot and mouth disease (Kamakawa et al., 2006).

Commercial movements of live animals are known to be a major pathway for disease transmission between domestic animal populations (Fevre et al., 2006). Therefore, a description of the pig value chain and the analysis of the network of commercial movements of pigs between farms would provide insights in the potential transmission dynamics of influenza and other swine diseases between farms. Analysis of this network would provide essential elements for the development of targeted surveillance strategies. Indeed, network analysis has been applied in preventive veterinary medicine since the early 2000s and used to assess the influence of the distribution of contacts between animal populations from different stages of the value chain - such as farms, markets or slaughterhouses - on the disease transmission dynamics (Martinez-Lopez et al., 2009b). The position of these premises in the network are assessed, allowing the identification of the premises at higher risk of becoming infected or of transmitting infection (Dube et al., 2011). In Vietnam, there is no record of live pig movements and only a limited number of publications addressing this topic are available. Some studies described the pork value chain in Vietnam focusing on the distribution of meat (Lapar et al., 2003; ILRI, 2014), and the cross-border movements of live pigs and other live-stock in South-East Asia (Cocks et al., 2009). To our knowledge, no study of the live pig trade network in Asian countries has been carried out, while a few studies have

been carried out in European countries and Canada since 2007 using commercial or governmental databases (Bigras-Poulin et al., 2007; Martinez-Lopez et al., 2009a; Noremark et al., 2011; Rautureau et al., 2012; Buttner et al., 2013; Dorjee et al., 2013; Smith et al., 2013; Thakur et al., 2014). In contrast, network analysis on poultry movements in Vietnam (Soares Magalhaes et al., 2010; Fournie et al., 2013), China (Martin et al., 2011; Soares Magalhaes et al., 2012), Cambodia (Van Kerkhove et al., 2009) and developing countries in other parts of the world such as Madagascar (Rasamoelina-Andriamanivo et al., 2014) was carried out. These studies conducted in Asia have pointed out the role of live bird markets in the dissemination of highly pathogenic avian influenza and provided useful insights for the surveillance and control of the disease in poultry.

The objective of this study was to describe farming and trading practices in the familial swine sector in the Red River Delta region (RRD) in northern Vietnam. Farmers in two communes in the RRD and traders involved in pig trades were interviewed. The geographical scale and the structure of the networks of contacts between farms resulting from pig trade were assessed. Hypotheses on the risk of disease transmission across the pig value chain were formulated for a comprehensive risk assessment that could be carried out in a separate future study. The identification of at risk trading behaviours of premises was useful in the design of risk-based surveillance protocols for the detection of infectious diseases such as swine influenza.

## 2. MATERIAL AND METHODS

### 2.1. Study area

The study was implemented in northern Vietnam in the RRD, where the capital city Hanoi is located. The RRD alone includes about a quarter of the human, pig and poultry populations of the country in just 6.4% of its area (GSO, 2011a,b, 2012). Hung Yen Province

was selected based on the following criteria: high density of pigs, economic importance of the pig production in the area and diversity of familial farming systems and trading practices. Following the same criteria as above, the study area was then narrowed down to two communes, Dinh Du (Com1) and Me So (Com2), within two adjacent districts, Van Lam (Dist1) and Van Giang (Dist2) located in the north of the province. These communes had similar characteristics such as the surface area (4.5 and 6.6 sq. km, respectively), human population (7100 and 9600 inhabitants) (GSO, 2014), number of villages (four and six) and number of familial farms (158 and 141) (May 2012 data). These two communes were especially chosen as they differed in the ratio of the large familial farms over small-medium familial farms: Com2 had 3.5 times more large farms and 1.4 times less small-medium farms compared to Com1. Indeed, the dynamic of farming intensification was different in the two communes, and this may have an impact on trading networks.

## **2.2. Selection of farmers and data collection**

Participatory interviews were carried out to collect preliminary data on the study areas and inform the design of the questionnaires. Different pig categories were mentioned. Pigs for slaughtering, later mentioned as slaughter pigs, included fattening pigs which were from 5-6 months old and incidentally cull pigs which were retired sows and boars. Weaners were about 2-month-old pigs that were fattened, and breeders included sows and boars for reproduction. The familial farms were categorized into small farms including *Nông hộ* (<100 fattening pigs per cycle and < 10 sows) and large farms including *Gia trai* and *Trang trai* ( $\geq 100$  fattening pigs per cycle or  $\geq 10$  sows). All the large farms were interviewed as their number was limited in each commune, whereas all the small farms from only one selected village in each commune were interviewed. Indeed, from the participatory interviews, it was estimated that the farming

and trading practices in each commune were homogeneous across the villages and the village with the highest number of small farms in each commune was selected.

Through questionnaire-based individual interviews, farmers were asked detailed information on the structure of the farm, their breeding practices and their purchases and sales of pigs from January 2011 to June 2012. Indeed, at least 1 year of trading period was chosen because of the duration of fattening pig production (around 6 months) and to account for the potential seasonal increase in pig production due to the Tết holiday, the Vietnamese New Year, occurring in January or February (during which consumption of meat increases). During the interviews, free recall and free choice approaches were used to identify all the persons they had traded with during the specified period of time; no a priori list of stakeholders was provided. Farmers with a larger number of trades were more likely to forget some of them, and also trades of smaller sizes may have been omitted more frequently. This recall bias may have led to an underestimation of the number of trades and pigs, especially for large farmers with many trades. The reports of trades between interviewed actors were checked for consistency later on.

## **2.3. Selection of traders and data collection**

The interviewed farmers mentioned different pig buyers and sellers. Within a snowball sampling approach, a sample of the actors involved in the trade of weaners and fattening pigs (the most important trades in terms of volume) was chosen for additional interview. This included companies and traders such as traders owning or working in slaughterhouses, middlemen buying and selling pigs directly between farmers, and middlemen buying pigs from farmers and selling them in the live pig market in Com1. For each category, the most mentioned actors with sufficient contact information were selected in priority

across Hung Yen, Hanoi and the surrounding provinces. Traders were asked general and detailed information on their activity, including questions about the number of trades and pigs exchanged from January 2011 to June 2012. Contrary to farmers, traders could not provide a detailed list of their trades as there were too many. Therefore, the total volume of trades and pigs traded over the study period were asked, and then, matrix scorings (Jost et al., 2010), a participatory method using proportions, were used to estimate the number of pigs traded and trades performed with each category of actor and location for both purchases and sales. All farmers' and traders' interviews were performed between June and September 2012.

## **2.4. Data processing**

Data were entered in an access database and were cleaned and analysed using R 3.1.1 (R Core Team, 2013). During farmers' interviews, potential inconsistencies between the answers provided to questions related to overall and specific trading activities were clarified. When important inconsistencies were noticed when cleaning the data set, the paper form was checked, and if the error was not resolved, the interviewees were contacted again by phone for clarification.

## **2.5. Data analysis**

### **2.5.1. Farm typology**

A farm typology taking into account both the size and the type of production was carried out, as these characteristics were considered to influence farmers' trading practices. A principal component analysis followed by a hierarchical clustering was conducted. Five variables were included: number of sows and number of boars present at the time of the visit, average number of fattening pigs produced per year in the farm, and number of weaners purchased and sold from 2011 to June 2012. The farm classes resulting from the typology were then used in the rest of the analysis.

### **2.5.2. Descriptive statistics**

First, we performed basic descriptive analyses of the farming and general trading practices. Then, a detailed description of the trades from farmers and traders was performed for the different pig production categories, that is slaughter pigs (fattenings and cull pigs), weaners and breeders. Fattening and cull pigs were not moved from farms to farms, but to slaughterhouses; these movements can be considered as dead ends for disease circulation. Greater focus was brought on the movements of pigs between farms, directly or through middlemen, which involved weaners and breeders. Cytoscape 3.2.0 (The Cytoscape Consortium, San Diego, CA, USA) was used to draw diagrams showing the movements of pigs between the different categories of actors (Smoot et al., 2011). Trades were qualified as transversal when performed between farms of equivalent sizes and as pyramidal when performed between farms of different size, for example from larger to smaller farms. For the graphs representing weaners and breeders trading from farmers in both communes, a bimodal approach was used, that is sales and purchases were treated and represented separately. Loops were eliminated by this process, and trades between interviewees were counted twice, once as sales and once as purchases to clarify the direction of trades, that is transversal or pyramidal.

### **2.5.3. Analysis of egocentric networks**

Commonly within an animal movement network, nodes represent individual premises part of the value chain, and the links between two nodes represent animal movements. The links are called arcs in a directed network where one of the nodes is the sender or seller and the other is the receiver or buyer (Dube et al., 2011). Egocentric networks were built for each of the 137 farms based on weaner and breeder trades using the R package igraph (Csardi and Nepusz, 2006). This analysis was chosen over the construction of one network in

each commune involving all interviewed and mentioned actors because about 90% of the trades were performed with non-interviewed actors and the networks would have been incomplete.

Each egocentric network was composed by nodes representing an interviewed farmer and his trading partners, mentioned as large or small farmers, companies, middlemen or market. The market was considered as one node with a unique location, although many middlemen would also be present in the market. Ideally, a node should be defined by a premise where pigs would stay at least a few hours like in slaughterhouses and markets, but some middlemen carried pigs directly from farms to farms. However, the farms of origin or destination were not known for middlemen trades, and therefore, it was not possible to replace these middlemen by arcs to farm nodes. The arcs represented the movements of weaners and/or breeders from nodes selling to nodes buying the pigs. For each interviewed farmer, two egocentric networks were built with arcs characterized by two different weights, being the number of trades and the number of pigs traded from January 2011 to June 2012. Thus, 274 egocentric networks were built. For each interviewed farmer, different values of degrees were calculated. The in- and out-degrees were the number of actors selling pigs to and purchasing pigs from this farmer, respectively. Similarly, the pig weighted in- and out-degrees were the total number of pigs a farmer purchased and sold, respectively. And finally, the trade- weighted in- and out-degrees were the total number of trades a farmer performed for purchase and sale. Clustering coefficients could not be calculated as most of the trades were performed with non-interviewed actors, and therefore in the majority of the egocentric networks, the links between the actors mentioned by the farmer were missing. Based on these six values of degrees, a network typology was carried out using the same methodology as for the farmtypology. The

correlation coefficients between the different variables were calculated, and when they were above 0.8, one of the two variables involved was excluded. As a result, the networks were grouped into different classes according to their degree values. Finally, a descriptive analysis of the networks and that of these different classes were performed in relation to the farm classes and other practices related to pig movements such as boar hiring.

#### ***2.5.4. Description of the geographical distribution of the trades***

First, the spatial distribution of the trades generated by farmers was compared at the province level between Com1 and Com2. Then, middleman trades were analysed to see whether they were likely to link different provinces through pig movements. Finally, the general movements of live pigs were described geographically for all pig categories based on farmers' and traders' interviews. A map of all these trades across Vietnam was drawn using ArcGIS 10.1 (ESRI, Redlands, CA, USA); for this purpose, the trades mentioned by the farmers with the interviewed traders were deleted as it was considered that these trades were included in the traders' transactions. For the trades performed with middlemen and markets, the trader origins were taken into account as the locations of the farms of origin or destination were unknown.

### **3. RESULTS**

#### **3.1. Study sample**

A total of 158 farmers were listed by the village veterinarians including all the large farms in both communes and the small farms in both selected villages. However, some pig farmers were not listed, some farmers had stopped their activity by the time the study started, and some were not available or refused the interview. Finally, 137 farmers were interviewed, including nine large and 40 small farmers in Com1, and 41 large and 47 small farms in Com2.

About 100 companies and traders in 11 provinces were identifiable by their name and were mentioned in 350 trading occasions. In 58 occasions, the information given by the interviewed farmer was not sufficient to identify the trading partner, 95% of these being slaughterhouses. Finally, a total of 11 slaughterhouses, 22 middlemen, and two companies were interviewed in five provinces, accounting for over 30% of all buyers and sellers mentioned and half of the trading occasions. The 22 interviewed middlemen included 17 middlemen exclusively trading weaners, with nine of them operating in weaner markets, three middlemen exclusively trading fattening pigs and two middlemen trading both weaners and fattenings. Only 10% of the companies mentioned were interviewed accounting for about 20% of the trading occasions with companies.

### **3.2. General production and trading practices**

Four classes of farms were obtained from the farm typology (Figure S1). Three classes represented a total of 41 large farms of different production types (farrow to finish and fattening), while the fourth class grouped 96 small farms without discriminating the production type. The class containing the 96 small farms was then divided into two production types equivalent to the production types obtained for the large farms. Thus, the 137 farmers were divided into five classes. A summary of the typology variables for each of the farm classes is presented in Table 1. A total of 62.8% of the farms were specialized in a finishing activity, that is fattening 2-month-old weaners until slaughtering, and were divided in two classes defined as large fattening farms (LF) and small fattening farms (SF). The three other classes were farrow-to-finish farms of different sizes with very large farms (VB), large farms (LB) and small farms (SB). Contrary to fattening farms, they did not need to purchase weaners due to their breeding practice. Among the LB farms, three had a

farrowing activity only. Although they were all familial farms, the VB farms had a very high number of pigs similar to some companies. Most of the large farms (VB, LB, LF) were found in Com2, while in Com1 and Com2, a similar number of small farms were interviewed in each village (SB, SF). These results show that the familial farms of different sizes have a specialized pig production which may influence their trading behaviours.

A descriptive analysis of trading practices was performed for the different pig categories in each commune (Table 2). Farmers in Com2 were trading more pigs (over 50 000 pigs traded) compared to Com1 (about 8000 pigs traded). This was especially true for weaners which were mainly purchased by local farmers in Com1, whereas in Com2, weaners were both purchased and sold in high quantities by the interviewees. Pig farming was the most important source of income for the majority of the interviewees, so they usually remembered the trades they had performed within the last 1.5 year. However, for the trades performed between interviewees, involving breeders and weaners, three quarters (37/49) of the links were only reported by one of the two interviewees involved in the exchange. The exchanges of larger numbers of pigs were usually better remembered as compared to smaller volume of trades. No other differences were found between the trades that were forgotten or remembered.

### **3.3. Trading practices for slaughter pigs**

The analysis of the sales of slaughter pigs among the farmers showed important differences between the two communes (Fig. 1). In Com1, farmers sold about three quarters of the pigs to traders slaughtering at the farm, while in Com2, a similar proportion of the pigs were sent to slaughterhouses. The farmers mentioned several categories of traders involved in fattening pig transactions. Two traders in collective slaughterhouses in Hanoi, nine local slaughterhouses and five middlemen in Hung Yen and adjacent provinces were interviewed,

**Table 1. Median (min-max) of the different variables  
for the different classes of swine farms identified**

Class	Sows	Boars	Weaners sold	Weaners purchased	Fattenings per year	No. of farms (Com1; Com2)
VB	185 (70-250)	5 (3-6)	1460 (0-6980)	0 (0-0)	1520 (600-2100)	4 (1; 3)
LB	26 (10-70)	1 (0-5)	112 (0-1320)	0 (0-0)	248 (0-700)	22 (5; 17)
SB	4 (1-12)	0 (0-0)	8 (0-61)	0 (0-34)	64 (20-160)	25 (12; 13)
LF	0 (0-4)	0 (0-0)	0 (0-0)	500 (280-970)	300 (140-900)	15 (0; 15)
SF	0 (0-13)	0 (0-0)	0 (0-0)	80 (10-253)	60 (15-300)	71 (31; 40)

*Note: VB, very large farrow-to-finish farms; LB, large farrow-to-finish farms; SB, small farrow-to-finish farms; LF, large fattening farms; SF, small fattening farms.*

**Table 2. General trade description per pig category  
in both communes over a 1.5-year period**

Type of trades in the two communes	Slaughter pigs		Weaners		Breeders	
	Pigs	Trades (p/t)	Pigs	Trades (p/t)	Pigs	Trades (p/t)
Total No in Com1	6086	363 (12)	1886	153 (12)	86	16 (5)
Purchases (%)	0	0	81.4	82.3	96.5	87.5
Sales (%)	100	100	13.1	11.8	0	0
Trades between interviewees (%)	0	0	5.5	5.9	3.5	12.5
Total No in Com2	24 243	1050 (26)	26 724	576 (40)	597	62 (8)
Purchases (%)	0	0	39.6	33.2	93.5	87.1
Sales (%)	98.6	98.4	53.0	55.0	0	0
Trades between interviewees (%)	1.4	1.6	7.4	11.8	6.5	12.9
Com2/Com1	4.0	2.9 (2.2)	14.2	3.8 (3.3)	6.9	3.9 (1.6)
Com2/Com1 adj. (a)	2.2	1.6	7.9	2.1	3.9	2.2

*Note: (p/t), Average number of pigs per trade. Bold values represent the total numbers of pigs and trades in each commune, and the ratios between the two communes. <sup>a</sup>Ratios adjusted by the number of farms in each commune.*

and their trading practices were described (Table S1). In the collective slaughterhouses, several independent traders were slaughtering pigs from many farms every night, mainly companies and large farms. In local slaughterhouses, pigs from only one or two farms were slaughtered per night, mostly from familial farms. Some seasonal variations were observed with an increase in the number of pigs slaughtered for a few days during the T  t holiday. However, this seasonal effect for pig trade was not acknowledged by farmers or middlemen and was probably limited. These practices will have a direct impact in terms of disease surveillance. Collective slaughterhouses

seem the most suitable for sampling pigs from a high number of farms. On the opposite side, pigs slaughtered at the farm will not be easily accessible for sampling.

### 3.4. Trading practices for weaners and breeders

The distribution of the trades for weaners and breeders according to the trading partner activities was represented in Fig. 2 for each pig category in each commune. In the two communes, mostly, fattening farms were responsible for the purchases of weaners. These pigs were mainly purchased indirectly through middlemen and the market (61.7%) by the small fattening farms in




Com1 and directly from other farmers (86.6%) by the large and small fattening farms in Com2. Farrow-to-finish farms in Com1 were selling only a few hundreds of weaners, while the one in Com2 was responsible for the sale of above 16 000 weaners. Overall, the large farms in Com1 were almost completely inactive in terms of weaner trades, while the three VB farms in Com2 accounted for 61.4% of weaners' sales in that commune.

The trades were qualified as pyramidal when the pigs were going from farms of larger sizes (companies, large farms) to farms of smaller sizes (large farms, small farms) and transversal when the pigs were exchanged between farms of same size. In Com1, the interviewed middlemen (n = 10) mentioned trading with many actors (Fig. 3a), while the traders interviewed in the market (n = 9) reported purchasing weaners mainly from small

farmers (Fig. 3b). Therefore, in Com1, the trades appeared to be mainly transversal between small farms, directly or through the market. In Com2, an important part of the trades was clearly pyramidal for both purchases (60.6%) and sales (50.1%). The companies interviewed reported selling 64.6% of their weaner production to familial farmers, confirming the strong pyramidal structure.

Regarding the breeder trades, only a few hundreds were purchased and only a few dozens were sold altogether. In both communes, most of these trades were performed directly between farms. The breeders were mainly purchased from companies, including breeding companies, and in general, the purchases were almost exclusively pyramidal trades. The sales were performed only locally in the same commune between familial farms.



**Fig. 1. Trade distribution for slaughter pigs across actor categories in each commune**


*Note: Percentages of pigs traded ( $n_{Com1} = 6086$ ,  $n_{Com2} = 24\ 243$ ); NA = buyer activity unknown, slaughter mixed = traders with different slaughtering activities.*

### 3.5. Analysis of egocentric networks

Ninetyeight egocentric networks in Com1 and 176 in Com2 were built from the interviewed farmers; they were describing the trades of weaners and breeders between actors. An additional 67 actors in Com1 and 139 actors in Com2 were mentioned by the farmers and composed the networks, accounting for a total of 97 and 261 arcs, respectively. The network sizes, that is the number of nodes, varied from one to 13 in Com1 and one to 22 in

Com2 with a median of 3.0 and 3.5, respectively. VB and LB farms in Com2 had larger networks compared to other farms in Com1 and Com2, confirming the higher trading dynamism of these farm classes in Com2.

For the network typology, five variables were kept after exclusion of the trade-weighted out-degree variable because of its high correlation coefficient with the pig-weighted out-degree (0.97) and the out-degree variables (0.82). Four classes were identified (Figure S2),



**Fig. 2. Trade distribution for weaners and breeders between actor categories in each commune**

*Note: (a1 and a2) trades of weaners in commune 1 and 2, respectively; (b1 and b2) trades of breeders in commune 1 and 2, respectively. Percentages of purchased weaners ( $n_{Com1} = 1,640$ ,  $n_{Com2} = 12,563$ ) and breeders ( $n_{Com1} = 86$  pigs,  $n_{Com2} = 597$ ) from the different actor categories and sold weaners ( $n_{Com1} = 351$ ,  $n_{Com2} = 16,131$ ) and breeders ( $n_{Com1} = 3$ ,  $n_{Com2} = 39$ ) to the different actor categories; trades between interviewees are counted twice as sales and purchases separately. Arrow widths are proportional to the number of pigs traded.*

and a summary of all the variables is available in Table 3. The first group was designated as isolated farms as they had null-to-low in-degree measures with null to medium out-degrees, that is they had limited trading interactions with other farms. The farms named as primary and secondary sinks had null-to-low out-degrees and high and medium in-degrees, respectively; therefore, they were receiving pigs but not redistributing. Finally, farms with high out-degrees were classified as sources, as they were at the origin of many sales; they also had medium in-degrees making them act as mediators for the movements of pigs, comparatively to the other farm classes. The companies were not included in the typology as a focus was performed on familial farms. However, they could also be considered as sources for the familial farms. The sources were VB and LB farms in Com2, while these same farm classes in Com1 were either sinks or isolated farms (Table 4). LF farms and about half of SF farms were primary sinks due to their important finishing activity. The remaining SF farms were mostly secondary sinks, that is with less important purchases. Most SB farms were isolated as they had

limited trades, probably due to their small size and self-sufficiency in terms of weaner supply with reproduction on the farm.

3.6. Boar hiring and its impact on the swine movement networks

Boar hiring was also mentioned as pig movements other than trades. About half of the farms with sows (42 farmers) reported hiring boar from other familial farmers. This practice was the most common in small farms and in Com1 where about 90% of farms with sows hired boars compared to about 50% in Com2 (Table S2). The network classes affected were the isolated farms and primary and secondary sinks. The degree variables for the farms hiring boars, originally based only on sales and purchases, were recalculated including and excluding the boar movements, to assess the impact of this practice on the classification of the farms (Table S3). In general, the number of pigs did not increase dramatically as only one boar was exchanged at a time, while the frequency of exchanges or trades increased consequently, connecting many originally isolated farms with others.




Fig. 3. Distribution of weaner trades generated by interviewed middlemen (a) and market traders (b)

Note: Percentages of weaners purchased and resold by independent middlemen ( $n_{\text{middlemen}} = 202,864$  pigs) and middlemen from the market in Com1 ( $n_{\text{market}} = 40,870$  pigs).

**Table 3. Median (min-max) of the different variables for the network classes identified**

Class	Out-degree	Trade-weighted out-degree	Pig-weighted out-degree	In-degree	Trade-weighted in-degree	Pig-weighted in-degree	Total no. of farms
Isolated farms	0 (0 - 4)	0 (0 - 5)	0 (0 - 239)	1 (0 - 1)	1 (0 - 2)	1 (0 - 40)	34
Primary sinks	0 (0 - 1)	0 (0 - 1)	0 (0 - 1)	3 (1 - 12)	5 (3 - 16)	176 (30 - 970)	48
Secondary sinks	0 (0 - 2)	0 (0 - 4)	0 (0 - 85)	2 (1 - 2)	3 (1 - 7)	50 (2 - 220)	42
Sources	6 (4 - 19)	18 (5 - 126)	690 (100 - 6980)	2 (1 - 3)	2 (1 - 5)	20 (3 - 117)	13

*Note: Degree measures were based on trades of weaners and breeders only.*

**Table 4. Proportions of farm classes pertain- ing to the network classes**

Class	VB	LB	SB	LF	SF
Isolated farms	-	31.8 (3; 4)	84.0 (12; 9)	-	8.4 (4; 2)
Primary sinks	-	-	-	100 (0; 15)	46.5 (12; 21)
Secondary sinks	50.0 (1; 1)	18.2 (2; 2)	16.0 (0; 4)	-	45.1 (15; 17)
Sources	50.0 (0; 2)	50.0 (0; 11)	-	-	-
Total no. of farms	4	22	25	15	71

*Note: Percentage and number of farms in commune 1 and in commune 2: 'Percentage (No in Com1; No in Com2)'.*

### 3.7. Spatial analysis of swine movements

Differences were observed in the spatial distribution of the farmer trades between sales and purchases, between communes, and between pig categories (Figure S3). In general, farmers in Com2 were trading with actors from multiple different provinces (4-7 provinces) compared to Com1(2-4 provinces). The largest proportion of pigs was traded with Hung Yen for slaughter pigs (87.0% and 56.5% for sales) and weaners (93.2% and 94.6% for sales, 75.2% and 36.2% for purchases) for Com1 and Com2. Most of the slaughter pig trades involved a limited number of provinces all located around Hung Yen. Weaners were traded between more provinces in the RRD and northern Vietnam. Breeder trades involved the highest number of provinces, including provinces from southern Vietnam, probably due to the high genetic quality of the breed offered in that region. Breeders were purchased in majority from Hanoi (51.2% in Com1 and 67.0% in Com2).

Thirteen provinces in the RRD and the north were mentioned by the traders for weaner trades and 13 provinces in the RRD, the north and the south for fattening trades (Figure S4 and S5). However, most of the traders were trading pigs within the RRD and especially Hung Yen and the three adjacent provinces. Among weaners and fattening middle men, three mentioned buying and selling in only one province (Hung Yen), while for 19 middlemen, the provinces used for purchases were not all the same than those mentioned for sales. This demonstrated that they were involved in pig movements between different provinces.


Overall, the trades generated by all interviewed traders and farmers for all pig categories involved the movement of a total of about 900 000 pigs across 22 provinces including Hung Yen over the 1.5-year study period (Fig. 4). They were distributed all over Vietnam, with a focus on the north and especially the RRD. Trades including the south were related to breeder purchases.

4. DISCUSSION

This study provided a detailed descriptive analysis of the farming systems and live pig movements in two communes of the RRD in northern Vietnam. The farm typology and the analysis of the farm networks provided important information for the identification of trading behaviours at risk for contagious disease spread and for the development of targeted surveillance strategies. Cost effective surveillance protocols are needed which allow the monitoring of diseases at a reasonable and therefore sustainable cost. Risk-based surveillance is the best approach and may have different goals (Cameron, 2012). Disease detection is a common goal, for example, for virus characterization, in the case of SIV in order to study the zoonotic potential of the viruses or for PRRS to develop vaccines matching the circulating strains. Another goal is to demonstrate freedom from disease; this is more often used in developed countries for trading purposes, while in Vietnam, most of the major swine diseases are endemic. Finally, early detection of a disease is often sought to contain its spread by implementing different

control measures (e.g. trade restrictions), for example in the case of a PRRS outbreak. Disease mitigation can also be performed by implementing preventive measures such as the increase in the general biosecurity level. Due to limited resources, farming systems at high risk of disease spread to an important number of other farms need to be targeted in priority. The design of risk-based surveillance protocols and control measures requires the identification of farm categories with a higher risk of disease introduction and maintenance on one hand and disease spread on the other hand, and the identification of suitable focal points for targeted sampling is also valuable.

Although the study sample was limited as it was designed to be exhaustive at the local level, our data analysis showed an important diversity of practices among familial farms. Moreover, as the study province is one of the most dynamic in terms of pig production in the RRD, it could be considered that most of the different trading behaviours found in this region were also found and documented in the study area. The study provided insights in the dynamic of local pig production and trades. However, as the study



**Fig. 4. Spatial distribution of the pig movements generated by interviewed farmers and traders**

*Note: The number of pigs purchased in and sold to a province is represented on the overall map of Vietnam and in a zoom of the RRD. The number of pigs exchanged within the same province is shown in a second zoom of the five provinces in the RRD where internal trades were mentioned*

focused on familial farms, the company sector was not thoroughly investigated and additional studies should focus on this sector to fully apprehend its role in terms of pig movements and risk in disease spread between sectors. It was pointed out that trades were sometimes forgotten by farmers, especially when small; therefore, the swine commercial exchanges may have been underestimated. In addition, because of the lack of detailed contact information in some instances, it was not possible to interview all of the trading partners mentioned by the interviewees and to perform an analysis of the complete trading network. Therefore, in future studies, we recommend that movement network analysis be carried out by asking farmers and traders to keep written records of all sales and purchases. In the current study, only traders near Hung Yen and Hanoi were interviewed, as most of the traders mentioned were located in that area. The geographical extent of the trades may therefore have been underestimated; however, many provinces were mentioned, suggesting that provinces in northern Vietnam are highly connected through the pig trade. As traders were not able to provide the exact number of pigs traded by actor category and location, only general trends were investigated using matrix scoring. This method does not provide precise numbers, but it provides reasonable estimates particularly of the relative proportions for the volume of pigs by actor and location. This analysis allowed a reasonable description of the farming systems, of their trading practices, and allowed farm classification according to the direction and volume of pig movements which could be interpreted in terms of risk of disease introduction and spread. The risk of disease introduction and spread in low biosecurity settings such as Vietnam familial farming is likely to be linked to trading behaviours and the corresponding animal and human movements.

Indeed, our typology of the familial farms performed within this study allowed going beyond the simple distinction between small and large farms. Our results showed that farms had a very specialized pig production, as

in industrialized countries, with farrow-to-finish and fattening farms of different sizes (Table 1). However, very few farms had an exclusive farrow-to-grower activity, and no farms had a nursery activity (pigs from three to 10 weeks of age) as it is described in Western countries (Noremark et al., 2011; Rautureau et al., 2012; Dorjee et al., 2013). On the overall farmers in Com1 had less dynamic trading practices as compared to those in Com2 (Table 2), where a higher level of intensification of the familial pig production was observed (with a higher number of large and very large farms). The trading practices were very different between the two communes and the different types of production. The structure of the trades differed between Com1 with mostly transversal trades (between farms of same size) and Com2 with mainly pyramidal trades (between farms of different size) (Fig. 2). Moreover, the pyramidal structure of the pig production had a double sense here, with pig movements going from larger to smaller farms, but also from farms with a farrowing activity to fattening farms. This type of trade structure is found in industrialized countries as well as described in the network analysis studies cited previously. Due to these trade structures, transmission of diseases through live pig movements is more likely to occur between farms of similar size through transversal trades and from larger farms towards smaller farms through pyramidal trades. In the absence of appropriate biosecurity measures, the companies could play an important role as sources of disease spread to the familial sector through the sales of breeders and weaners. Specific types of production also had specific trading practices and thus different risk of disease spread. In terms of disease transmission risk through trade, sources are generally considered at high risk of disease spread to other farms, and sinks have a high risk of receiving the disease. Here, the farms defined as sinks were fattening farms with important purchases of weaners; they are potentially at higher risk of disease introduction due to pig movements, but

probably have a lower chance of spreading the disease through trade due to their limited number of sales to other farmers, as pigs are mainly sent to slaughterhouse (Tables 3 and 4). The source farms, which were large farrow-to-finish farms with moderate purchases (replacement of breeders) and important sales to other farmers (mainly weaners and some breeders), would have a non-negligible risk of disease introduction and a high risk of spreading the disease to other herds. On the opposite, the small farrow-to-finish farms in both commune and the large ones in Com1 were considered as isolated farms and would therefore play a limited role in disease transmission risk through pig trading networks. No farms showed both high in- and out-degrees, and therefore they are not at high risk of disease introduction and spread (act as a hub), as it was described for nursery farms in a study in Canada (Dorjee et al., 2013).

Therefore, one could consider that farms in Com1 would be less at risk for disease spread, having a more limited number of trades. However, other at-risk practices, also shared with some farms in Com2, were identified through the study. Firstly, middlemen and market were more frequently involved in Com1 swine purchase activities which increase their risk of disease introduction (Figs 1 and 2). Indeed, middlemen often kept traded pigs in their house where they also sometimes raise pigs. In the market, pigs would stay for a few hours next to other pigs with different origins, sometimes being transported back and forth to the trader's house and potentially to other markets if unsold.

The risk of transmission by fomites and people during these trades could also be important. Farmers in Com2 preferred buying pigs directly from other farmers so they could check the health status of the herd. Secondly, in Com1, more farmers were hiring boars for reproduction, being more popular among small farmers in general (Figures S4 and S5). This practice may increase the role in disease transmission of farms considered as isolated

regarding the other types of trades, through boar infection or fomite contamination.

The spatial distribution of the trades observed in this study highlighted the geographical extent of the pig movements, although most of the pigs were usually traded within the RRD region (Fig. 4). Overall, over 20 provinces in all Vietnam, that is about a third of the provinces in the country, were mentioned for trades described in this study which involved 173 actors interviewed in a limited geographical area. Based on these data, swine diseases are likely to spread easily and quickly within the RRD and northern Vietnam as controls and health certificate requirements are not well regulated and because of the occurrence of asymptomatic diseases such as swine influenza.

According to the study results, fattening farms with high numbers of pigs purchased (i.e. sink farms) should be targeted in the design of a risk-based surveillance for disease detection and virus isolation for genetic characterization. Indeed, targeting these farms would increase the sensitivity of surveillance compared with a randomized sample including all farms. This is probably true for fattening companies with low biosecurity as well. The large farrow-to-finish farms buying breeders and selling weaners seemed the most at risk for disease spread. They should be targeted along with companies for implementing disease prevention and control measures because of their high potential to spread diseases to other farms. This could be performed by improving the general biosecurity level of these farms, and also by the early detection of diseases in these swine herds before it spreads down to the chain. As these farms are large and with constant renewal of the susceptible pig population, disease persistence might be high; they could also be considered in a risk-based surveillance design for disease detection, together with the fattening farms. Studying the disease transmission dynamics within these farms may be of interest to support the decision to include them in risk-based surveillance

protocols. Similarly, larger fattening farms may show a higher persistence of viruses and may be more of interest compared to smaller farms. Developing risk-based surveillance protocols in areas like Com1 may be more difficult as no farms with high risk were identified, and few large farms were present. Although other risky behaviours were more frequent in this commune like boar hiring, the chances of isolating viruses might be low as the window of infection in such small pig herd would be short, and therefore, the surveillance programme might not be cost-effective.

Finally, this study also allowed the identification and description of focal points that concentrate pigs from many farms. Sampling in these places would require fewer resources and would be more effective than sampling in individual farms. Local slaughterhouses were not considered as an efficient location for risk-based surveillance as they usually hosted pigs from only a few farms in a given day (Table S1). On the contrary, collective slaughterhouses represented a very promising candidate for risk-based surveillance design for disease detection because pigs slaughtered in a same night had come from many farms and many provinces. Pigs were usually transported from northern provinces within the same day and held at the slaughterhouse for a few hours up to 36 h before slaughter. Pigs from several farms were often mixed in the same pens, and this may facilitate cross-infection of non-immune animals and amplification of virus within this setting. Pigs in collective slaughterhouses came mostly from companies and secondarily from large familial farms. The other type of focal point identified was the weaner market. As the pigs present originated mostly from small familial farms in the RRD, this location was considered as interesting to cover this farming sector. However, this type of markets was not frequent in the region, and only three were identified with a geographical coverage much smaller compared to collective slaughterhouses. It was reported that sick pigs were often sold for slaughter, but in the case of the market, it

would be likely that middlemen would sell only healthy-looking pigs to farmers, although some diseases like influenza may be asymptomatic. The age of the pigs sampled would also have an impact, with weaners being young pigs which might still be partially protected by maternal antibodies and fattening pigs being older pigs that may have been infected earlier in life and not be shedding virus at the time of slaughter. In the example of swine influenza viruses, to date, virus isolation in Vietnam has only been reported in the company or industrial sector (Ngo et al., 2012), although there is serological evidence for influenza circulation in the familial sector (Trevennec et al., 2012). This study highlighted the challenges brought by the structure of the pig value chain which would explain this low detection rate of swine influenza in the familial sector and provide new information on how to overcome such challenges for cost-effective surveillance design of infectious diseases in swine.

## 5. CONCLUSIONS

This study has provided empirical data on the organization of the pig value chain in Hung Yen Province in the RRD. Indeed, the analysis of animal movements represents a challenge in developing countries where systematic record keeping is not well established. This study allowed us to formulate hypotheses on disease transmission between farming systems and geographical locations, providing critical information for the design of risk-based surveillance protocols. Pilot trials of these protocols are currently ongoing to identify the most cost-effective protocols for swine influenza surveillance in Vietnam.

## ACKNOWLEDGEMENTS

We thank staff members of the Hanoi University of Agriculture and CIRAD colleagues in Hanoi for providing useful information on the swine industry in Vietnam and helping in the data collection in the field. Also, we thank the province, district, commune



and village local authorities in Hung Yen Province for providing assistance in the data collection, as well as all the participants, that is veterinarians, farmers and traders, for their contribution. The project was funded in part by CIRAD, by the Area of Excellence Scheme of the University Grants Committee [AoE/ M-12/06] of Hong Kong, the Harvard Center for Communicable Disease Dynamics from the National Institute of General Medical Sciences (grant no. U54 GM088558) and the National Institutes of Health (NIAID contract HHSN272201400006C).

## REFERENCES

- Bigras-Poulin, M., K. Barfod, S. Mortensen, and M. Greiner, 2007: Relationship of trade patterns of the Danish swine industry animal movements network to potential disease spread. *Prev. Vet. Med.* 80, 143-165.
- Buttner, K., J. Krieter, A. Traulsen, and I. Traulsen, 2013: Static network analysis of a pork supply chain in Northern Germany - characterisation of the potential spread of infectious diseases via animal movements. *Prev. Vet. Med.* 110, 418-428.
- Cameron, A. R., 2012: The consequences of risk-based surveillance: developing output-based standards for surveillance to demonstrate freedom from disease. *Prev. Vet. Med.* 105, 280-286.
- Cocks, P., R. Abila, A. Bouchot, C. Benigno, S. Morzaria, P. Inthavong, N. V. Long, N. Bourgeois-Luthi, A. Scoizet, and S. Sieng, 2009: FAO ADB and OIE SEAFMD Study on Cross-Border movement and market chains of large ruminants and pigs in the Greater Mekong Sub-Region. FAO ADB and OIE SEAFMD, Thailand.
- Coker, R. J., B. M. Hunter, J. W. Rudge, M. Liverani, and P. Hanvoravongchai, 2011: Emerging infectious diseases in southeast Asia: regional challenges to control. *Lancet* 377, 599-609.
- Csardi, G., and T. Nepusz, 2006: The igraph software package for complex network research. *InterJ. Complex Syst.* 1695. Available at <http://igraph.sf.net> (accessed December 04, 2014).
- Dorjee, S., C. W. Revie, Z. Poljak, W. B. McNab, and J. Sanchez, 2013: Network analysis of swine shipments in Ontario, Canada, to support disease spread modelling and risk-based disease management. *Prev. Vet. Med.* 112, 118-127.
- Dube, C., C. Ribble, D. Kelton, and B. McNab, 2011: Introduction to network analysis and its implications for animal disease modelling. *Rev. Sci. Tech.* 30, 425-436.
- FAO, 2014: Food and Agriculture Organization of the United Nations. Global Livestock Production and Health Atlas. Available at <http://kids.fao.org/glipha/#> (accessed 24 October 2014).
- Fevre, E. M., B. M. Bronsvoort, K. A. Hamilton, and S. Cleaveland, 2006: Animal movements and the spread of infectious diseases. *Trends Microbiol.* 14, 125-131.
- Fisher, H., and J. Gordon, 2008: Breeding and Feeding Pigs in Vietnam: Assessment of Capacity Building and an Update on Impacts. ACIAR Impact Assessment Series Report. ACIAR, Australia. 52, 56 pp.
- Fournie, G., J. Guitian, S. Desvaux, V. C. Cuong, H. Dung do, D. U. Pfeiffer, P. Mangtani, and A. C. Ghani, 2013: Interventions for avian influenza A (H5N1) risk management in live bird market networks. *Proc. Natl Acad. Sci. USA*, 110, 9177-9182.
- GSO, 2011a: General Statistics Office of Vietnam. Number of pigs by province in 2011. Available at <http://www.gso.gov.vn/> (accessed October 24, 2014).
- GSO, 2011b: General Statistics Office of Vietnam. Number of poultry by province in 2011. Available at: <http://www.gso.gov.vn/> (accessed October 24, 2014).
- GSO, 2012: General Statistics Office of Vietnam. Area, population and population density in 2012 by province. Available at: <http://www.gso.gov.vn/> (accessed October 24, 2014).
- GSO, 2014: Vietnamese government information portal. Available at: <http://gis.chinhphu.vn/vbdmap.aspx#> (accessed October 24, 2014).
- ILRI, 2014: Improving the competitiveness of pig producers in an adjusting Vietnam market. Available at: <http://www.viet-pigs.com.vn/> (accessed October 23, 2014).
- Jones, K. E., N. G. Patel, M. A. Levy, A. Storeygard, D. Balk, J. L. Gittleman, and P. Daszak, 2008: Global trends in emerging infectious diseases. *Nature* 451, 990-993.
- Jones, B. A., D. Grace, R. Kock, S. Alonso, J. Rushton, M. Y. Said, D. McKeever, F. Mutua, J. Young, J. McDermott, and D. U. Pfeiffer, 2013: Zoonosis emergence linked to agricultural intensification and environmental change. *Proc. Natl Acad. Sci. USA* 110, 8399-8404.

- Jost, C. C., S. Nzietchueng, S. Kihu, B. Bett, G. Njogu, E. S. Swai, and J. C. Mariner, 2010: Epidemiological assessment of the Rift Valley fever outbreak in Kenya and Tanzania in 2006 and 2007. *Am. J. Trop. Med. Hyg.* 83, 65-72.
- Kamakawa, A., T. V. Ho, and S. Yamada, 2006: Epidemiological survey of viral diseases of pigs in the Mekong delta of Vietnam between 1999 and 2003. *Vet. Microbiol.* 118, 47-56.
- Lapar, L., T. B. Vu, and S. Ehui, 2003: Identifying barriers to entry to livestock input and output markets in Southeast Asia. *FAO Livestock sector report*, FAO, Vietnam.
- Martin, V., X. Zhou, E. Marshall, B. Jia, G. Fusheng, M. A. Francodixon, N. Dehaan, D. U. Pfeiffer, R. J. Soares Magalhaes, and M. Gilbert, 2011: Risk-based surveillance for avian influenza control along poultry market chains in South China: the value of social network analysis. *Prev. Vet. Med.* 102, 196-205.
- Martinez-Lopez, B., A. M. Perez, and J. M. Sanchez-Vizcaino, 2009a: Combined application of social network and cluster detection analyses for temporal-spatial characterization of animal movements in Salamanca, Spain. *Prev. Vet. Med.* 91, 29-38.
- Martinez-Lopez, B., A. M. Perez, and J. M. Sanchez-Vizcaino, 2009b: Social network analysis. Review of general concepts and use in preventive veterinary medicine. *Transbound Emerg. Dis.* 56, 109-120.
- Ngo, L. T., Y. Hiromoto, V. P. Pham, H. T. Le, H. T. Nguyen, V. T. Le, N. Takemae, and T. Saito, 2012: Isolation of novel triple-reassortant swine H3N2 influenza viruses possessing the hemagglutinin and neuraminidase genes of a seasonal influenza virus in Vietnam in 2010. *Influenza Other Respir. Viruses* 6, 6-10.
- Noremark, M., N. Hakansson, S. S. Lewerin, A. Lindberg, and A. Jonsson, 2011: Network analysis of cattle and pig movements in Sweden: measures relevant for disease control and risk based surveillance. *Prev. Vet. Med.* 99, 78-90.
- R Core Team, 2013: R: A Language and Environment for Statistical Computing. R Foundation for Statistical Computing, Vienna, Austria. Available at: <http://www.R-project.org/> (accessed November 26, 2014).
- Rasamoelina-Andriamanivo, H., R. Duboz, R. Lancelot, O. F. Maminiaina, M. Jourdan, T. M. Rakotondramaro, S. N. Rakotonjanahary, R. S. de Almeida, Rakotondravao, B. Durand, and V. Chevalier, 2014: Description and analysis of the poultry trading network in the Lake Alaotra region, Madagascar: implications for the surveillance and control of Newcastle disease. *Acta Trop.*, 135, 10-18.
- Rautureau, S., B. Dufour, and B. Durand, 2012: Structural vulnerability of the French swine industry trade network to the spread of infectious diseases. *Animal*, 6, 1152-1162.
- Smith, T. C., A. L. Harper, R. Nair, S. E. Wardyn, B. M. Hanson, D. D. Ferguson, and A. E. Dressler, 2011: Emerging swine zoonoses. *Vector Borne Zoonotic Dis.* 11, 1225-1234.
- Smith, R. P., A. J. Cook, and R. M. Christley, 2013: Descriptive and social network analysis of pig transport data recorded by quality assured pig farms in the UK. *Prev. Vet. Med.* 108, 167-177.
- Smoot, M. E., K. Ono, J. Ruschinski, P. L. Wang, and T. Ideker, 2011: Cytoscape 2.8: new features for data integration and network visualization. *Bioinformatics* 27, 431-432.
- Soares Magalhaes, R. J., A. Ortiz-Pelaez, K. L. Thi, Q. H. Dinh, J. Otte, and D. U. Pfeiffer, 2010: Associations between attributes of live poultry trade and HPAI H5N1 outbreaks: a descriptive and network analysis study in northern Vietnam. *BMC Vet. Res.* 6, 10.
- Soares Magalhaes, R. J., X. Zhou, B. Jia, F. Guo, D. U. Pfeiffer, and V. Martin, 2012: Live poultry trade in Southern China provinces and HPAIV H5N1 infection in humans and poultry: the role of Chinese New Year festivities. *PLoS One* 7, e49712.
- Thakur, K. K., C. W. Revie, D. Hurnik, Z. Poljak, and J. Sanchez, 2014: Analysis of swine movement in four Canadian regions: network structure and implications for disease spread. *Transbound Emerg. Dis.* doi: 10.1111/tbed.12225.
- Trevennec, K., L. Leger, F. Lyazrhi, E. Baudon, C. Y. Cheung, F. Roger, M. Peiris, and J. M. Garcia, 2012: Transmission of pandemic influenza H1N1 (2009) in Vietnamese swine in 2009-2010. *Influenza Other Respir. Viruses* 6, 348-357.
- Van Kerkhove, M. D., S. Vong, J. Guitian, D. Holl, P. Mangtani, S. San, and A. C. Ghani, 2009: Poultry movement networks in Cambodia: implications for surveillance and control of highly pathogenic avian influenza (HPAI/H5N1). *Vaccine* 27, 6345-6352.
- Vijaykrishna, D., G. J. Smith, O. G. Pybus, H. Zhu, S. Bhatt, L. L. Poon, S. Riley, J. Bahl, S. K. Ma, C. L. Cheung, R. A. Perera, H. Chen, K. F. Shortridge, R. J. Webby, R. G. Webster, Y. Guan, and J. S. Peiris, 2011: Long-term evolution and transmission dynamics of swine influenza A virus. *Nature* 473, 519-522.